



## Spotted Knapweed

**NATIVE RANGE:** Central Europe, central Russia, Caucasia, and western Siberia.


**DESCRIPTION:** Spotted knapweed, *Centaurea biebersteinii* DC. is a biennial or short-lived perennial member of the Sunflower (Asteraceae) family. Spotted knapweed typically forms a basal rosette of leaves in its first year and flowers in subsequent years. Rosette leaves are approximately 8 inches long by 2 inches wide, borne on short stalks, and deeply lobed once or twice on both sides of the center vein, with lobes oblong and wider toward the tip. The taproot is stout and deep. Flowering

stems are erect, 8 to 50 inches tall, branched above the middle, and sparsely to densely hairy. Stem leaves alternate along the stem, are unstalked, and may be slightly lobed, or linear and unlobed. Leaf size decreases towards the tip of the stem. Flowers are purple to pink, rarely white, with 25 to 35 flowers per head. Plants bloom from June to October, and flower heads usually remain on the plant. Flower heads are oblong or oval shaped, ¼ inch wide and ½ inch across, and are single or borne in clusters of two or three at the branch ends. Leaf like bracts surrounding the base of the flower head are oval and yellow green, becoming brown near the base. The margins of these bracts have a soft spine like fringe, with the center spine being shorter than the lateral spines. The brown, oval seeds are 1/16 to 1/8 inch long, with pale longitudinal lines and a short fringe on one end.

**ECOLOGICAL THREAT:** Spotted knapweed infests barrens, fields, forests, prairies, meadows, pastures, and rangelands. It out-competes native plant species, reduces native plant and animal biodiversity, and decreases forage production for livestock and wildlife. Spotted knapweed may degrade soil and water resources by increasing erosion, surface runoff, and stream sedimentation.

**DISTRIBUTION IN THE UNITED STATES:** (see green areas on map)

**HABITAT IN THE UNITED STATES:** Spotted knapweed is found at elevations up to and over 10,000 feet and in precipitation zones receiving 8 to 80 inches of rain annually. Spotted knapweed prefers well-drained, light-textured soils that receive summer rainfall, including open forests dominated by ponderosa pine and Douglas fir, and prairie habitats dominated by Idaho fescue, bluebunch wheatgrass, and needle-and-thread grass. Disturbance allows for rapid establishment and spread; however, spotted knapweed is capable of invading well managed rangelands.



### MANAGEMENT & CONTROL:

**Manual and Mechanical:** Small infestations of spotted knapweed can be controlled by persistent hand-pulling done prior to seed set. Gloves should be worn because of the possibility of skin irritation. Because spotted knapweed can regrow from the base, care must be taken to remove the entire crown and taproot.

**Biological control:** Two species of seed head flies, *Urophora affinis* and *U. quadrifasciata*, are well-established on spotted knapweed. The larvae of these species reduce seed production by as much as 50% by feeding on spotted knapweed seed heads and causing the plant to form galls. Three moth species (*Agapeta zoegana*, *Pelochrista medullana*, and *Pterolonche inspersa*) and a weevil (*Cyphocleonus achates*) that feed on spotted knapweed roots have also been released. Biological control agents may be more effective when combined with other control methods such as herbicides, grazing, and revegetation with desirable, competitive plants.

**Chemical:** Picloram is a persistent herbicide used to control knapweed for three to five years when applied at 0.25 lb/acre at any stage of plant growth; or with clopyralid (0.24 lb/acre) or clopyralid (0.2 lb/acre) plus 2,4-D (1 lb./acre) applied during bolt or bud growth stage. In the absence of desirable native grasses, longevity of control may be increased by revegetating with competitive grasses and forbs.

**Other methods:** Long-term grazing by sheep and goats has been found to control spotted knapweed. Burning, cultivation, and fertilization typically are not effective on spotted knapweed unless combined with other methods of control.

**References:** [www.invasivespecies](http://www.invasivespecies), [www.nps.gov](http://www.nps.gov)